

QCD ORIGINATED DYNAMICAL SYMMETRY FOR HADRONS

Djordje Šijački

Institute of Physics, P.O.B. 57, Belgrade, Yugoslavia

We extend previous work on the IR regime approximation of QCD in which the dominant contribution comes from a dressed two-gluon effective metric-like field $G_{\mu\nu} = g_{ab}B_\mu^a B_\nu^b$ (g_{ab} a color $SU(3)$ metric). The ensuring effective theory is represented by a pseudo-diffeomorphisms gauge theory. The second-quantized $G_{\mu\nu}$ field, together with the Lorentz generators close on the $\overline{SL}(4, R)$ algebra. This algebra represents a spectrum generating algebra for the set of hadron states of a given flavor - hadronic "manifolds" transforming w.r.t. $\overline{SL}(4, R)$ (infinite-dimensional) unitary irreducible representations. The equations of motion for the effective pseudo-gravity are derived from a quadratic action describing Riemannian pseudo-gravity in the presence of shear ($\overline{SL}(4, R)$ covariant) hadronic matter currents. These equations yield p^{-4} propagators, i.e. a linearly rising confining potential $H(r) \sim r$, as well as linear $J \sim m^2$ Regge trajectories. The $\overline{SL}(4, R)$ symmetry based dynamical theory for the QCD IR region is applied to hadron resonances. All presently known meson and baryon resonances are successfully accommodated and various missing states predicted.

$sl(4,R)$ SPECTRUM GENERATING ALGEBRA

One of the main challenges in Particle Physics is the understanding and/or classification of quite a large number of presently known hadronic resonances. Here we are faced with an intriguing situation: In the "horizontal" direction one has flavor symmetries and rather powerful quantitative techniques with practically none understanding of the corresponding underlying fundamental interaction. As for the "vertical" direction (fixed flavor content), the basic interaction is given by the presently widely accepted Quantum Chromodynamics (QCD) theory, however the non-perturbative features of QCD have made it difficult to apply the theory exactly. Quite a number of models and approaches have been proposed so far with different degree of success. We believe that the merits of the approach described in this article are both the fact that our starting point is QCD itself and that the predictions fit very well with experiment.

If the hadron lowest ground states are colorless (our assumption) and in the approximation of an external QCD potential, the hadron spectrum above these levels will be generated by color-singlet quanta, whether made of dressed two-gluon configurations, three-gluons, Every possible configuration will appear. No matter what the mechanism responsible for a given flavor state, the next vibrational, rotational or pulsed excitation corresponds to the "addition" of one such collective color-singlet multigluon quantum superposition. In the fully relativistic QCD theory, these contributions have to come from summations of appropriate Feynman diagrams, in which dressed n -gluon configurations are exchanged. We rearrange the sum by lumping together contributions from n -gluon irreducible parts, $n = 2, 3, \dots, \infty$ and with the same Lorentz quantum numbers. The simplest such system will have the quantum numbers of di-gluon, i.e. $n = 2$. The color singlet external field can thus be constructed from the QCD gluon field as a sum (g_{ab} is the $SU(3)$ metric, d_{abc} is the totally symmetric $8 \otimes 8 \otimes 8 \rightarrow 1$ coefficient)

$$g_{ab}B_\mu^a B_\nu^b \oplus d_{abc}B_\mu^a B_\nu^b B_\sigma^c \oplus \dots \quad (1)$$

In the above, B_μ^a is the dressed gluon field. It will be useful for the applications to separate the "flat connection" N_μ^a , i.e. the zero-mode of the field. Writing for the curvature or field strength $F_{\mu\nu}^a = \partial_\mu B_\nu^a - \partial_\nu B_\mu^a - if_{bc}^a B_\mu^b B_\nu^c$, we define

$$B_\mu^a = N_\mu^a + A_\mu^a, \quad (2)$$

so that $F(N) = 0$, i.e. $\partial_\mu N_\nu^a - \partial_\nu N_\mu^a = if_{bc}^a N_\mu^b N_\nu^c$. We can now rewrite the two-gluon ("di-gluon") configuration as [1]

$$G_{\mu\nu}(x) = g_{ab}B_\mu^a B_\nu^b \quad (3)$$

which looks very much like a spacetime metric ("pseudo-metric"). *We assume that (3) is the dominating configuration in the excitation systematics.* The separation of the flat part of B_μ^a in Eq. (2) reproduces the separation of a tetrad $e_\mu^a(x) = \delta_\mu^a + f_\mu^a(x)$ into the flat background piece and the quantum gravitational contribution. As a result, $G_{\mu\nu}(x)$ itself can be separated similarly. We discuss below the precise meaning of "pseudo-metric", "pseudo-gravity", "gauging",

The di-gluon external field $G_{\mu\nu}(x)$ transforms under Lorentz transformations as a (reducible) second-rank symmetric tensor field, with Abelian components, i.e. $[G_{\mu\nu}, G_{\rho\sigma}] = 0$. Algebraically, $G_{\mu\nu}$ and the Lorentz generators form the algebra of $T_{10} \circledast SO(1, 3)$, an inhomogeneous Lorentz group with tensor "translations" (the symbol $\circledast$ denotes a semi-direct product). This is a classical relativistic algebra. For the quantum case, when the gluon field is expanded in creation and annihilation operators, we can write [1]

$$G_{\mu\nu} = T_{\mu\nu} + U_{\mu\nu},$$

where the quadrupolar excitation-rate is given by

$$T_{\mu\nu} = g_{ab} \int d\tilde{k} [\alpha_\mu^{a\dagger}(k) \alpha_\nu^{b\dagger}(k) \exp(2ikx) + \alpha_\mu^a(k) \alpha_\nu^b(k) \exp(-2ikx)], \quad (4)$$

whereas

$$U_{\mu\nu} = \eta_{ab} \int d\tilde{k} [\alpha_\mu^{a\dagger}(k) \alpha_\nu^b(k) + \alpha_\mu^a(k) \alpha_\nu^{b\dagger}(k)], \quad (5)$$

We have made use of the canonical transformation $[\alpha_\mu^a(k) + \frac{1}{2}N_\mu^a \exp(ikx)] \rightarrow \alpha_\mu^a(k)$, $[\alpha_\mu^{a\dagger}(k) + \frac{1}{2}N_\mu^a \exp(-ikx)] \rightarrow \alpha_\mu^{a\dagger}(k)$. Using $[\alpha_\mu^a(k), \alpha_\nu^{b\dagger}(k')] = \delta^{ab} \delta_{\mu\nu} \delta(k - k')$, one verifies that the operators $T_{\mu\nu}$ and $U_{\mu\nu}$ together with the operators

$$S_{\mu\nu} = \eta_{ab} \int d\tilde{k} [\alpha_\mu^{a\dagger}(k) \alpha_\nu^b(k) - \alpha_\mu^a(k) \alpha_\nu^{b\dagger}(k)], \quad (6)$$

close respectively on the $gl(4, R)$ and $u(1, 3)$ algebras. Note that the largest (linearly realized) algebra with generators quadratic in the $\alpha_\mu^\dagger, \alpha_\mu$ operators is $sp(1, 3, R)$, where the notation "1, 3" implies a definition over Minkowski space.

$$sp(1, 3, R) \supset \left\{ \begin{array}{ccc} u(1, 3) & \supset & su(1, 3) \\ gl(4, R) & \supset & sl(4, R) \\ t_{10} \circledast so(1, 3) & \supset & t_9 \circledast so(1, 3) \end{array} \right\} \supset so(1, 3) \quad (7)$$

The $gl(4, R)$ algebra represents a Spectrum Generating Algebra for the set of hadron states of a given flavor [2,3].

We now return to the expansion in Eq. (1). The $sl(4, R)$ is generated by di-gluon. What about three-gluon and n-gluon exchanges? The corresponding algebras do not close and generate the full Ogievetsky algebra of the diffeomorphisms in Minkowski space, the four-dimensional analog of the Virasoro algebra (the algebra of diffeomorphisms on the circle).

Had we considered the entire (infinite) sequence when writing Eq. (1), we would have generated this $diff(4, R)$. The maximal linear subalgebra of $diff(4, R)$ is $gl(4, R)$. The remaining generators (i.e. $diff(4, R)/gl(4, R)$) can be explicitly realized in terms of the $gl(4, R)$ generators [4], both for tensors and for spinors. In our case, this would involve functions as matrix elements of the representation of our generators $T_{\mu\nu}$ and $S_{\mu\nu}$ in Eqs. (4) and (6). But $diff(1, 3, R)$ can also be represented linearly. It will then involve infinite,

ever more massive, repetitions of the representations of $sl(4, R)$. *In either way, we find that using $sl(4, R)$ takes care of the entire sequence in Eq. (1).*

The $SL(4, R)$ group preserves the four-dimensional measure, a geometric realization of confinement as a dynamical four-volume-preserving rotation-deformation-vibration pulsation mechanism. The $SL(4, R)$ algebra generators $Q_{\mu\nu}$, $\mu\nu = 0, 1, 2, 3$ satisfy the following commutation relations

$$[Q_{\mu\nu}, Q_{\rho\sigma}] = i\eta_{\nu\rho}Q_{\mu\sigma} - i\eta_{\mu\sigma}Q_{\rho\nu}, \quad (8)$$

where $\eta_{\mu\nu}$ is the Minkowski metric. The antisymmetric part of $Q_{\mu\nu}$, i.e. $M_{\mu\nu} = Q_{[\mu\nu]}$ generates the metric-preserving Lorentz group $SO(1, 3)$, while the remaining nine symmetric operators, i.e. $T_{\mu\nu} = Q_{(\mu\nu)}$ generate the relativistic four-volume-preserving transformations. In the following we will make use of the physically relevant component operators [5]:

$J_i = \frac{1}{2}\epsilon_{ijk}M_{jk}$	angular momentum,
$K_i = M_{0i}$	Lorentz boosts,
T_{ij}	shears (Regge sequence generating),
$N_i = T_{0i}$	
T_{00}	"time scaling"

The relevant subgroups are $SO(3)$ spatial rotations, generated by J_i , $SO(4)$ the maximal compact subgroup, generated by J_i and N_i , $SO(1, 3)$ the Lorentz subgroup, generated by J_i and K_i , $SL(3, R)$ the three-volume preserving subgroup, generated by J_i and T_{ij} , and R^+ the non-compact Abelian subgroup generated by T_{00} .

The maximal compact subgroup $SO(4) \simeq [SU(2) \otimes SU(2)]/Z_2$ is generated by the two commuting compact vector operators $J_i^{(1)} = \frac{1}{2}(J_i + N_i)$, $J_i^{(2)} = \frac{1}{2}(J_i - N_i)$, $i = 1, 2, 3$. The remaining nine (noncompact) operators transform with respect to $SO(4)$ as the components of the (1,1) irreducible tensor operator $Z_{\alpha\beta}$, $\alpha, \beta = 0, \pm 1$.

The inhomogeneous versions of the algebras in Eq. (7), i.e. their semidirect product with the translations t_4 , are relevant to the Hilbert space spectrum of states. In the case of $u(1, 3)$ in Eq. (7), when selecting a time-like vector (for massive states), the stability subgroup is the compact $u(3)$ with finite representations – as against the noncompact $gl(3, R)$ for $sl(4, R)$. This fits with the situation in nuclei, where symmetries such as the $u(6)$ of the IBM model [6] are physically realized over pairs of "valency" nucleons outside of closed shells. There is a finite number of such pairs, and the excitations thus have to fit within finite representations.

PSEUDO-GRAVITY DYNAMICS

The di-gluon effective gravity-like "potential" $G_{\mu\nu}$ acts as a "pseudo-metric" field, (passively) gauging effective "pseudo-diffeomorphisms", just as is done by the physical Einstein metric field for the "true" diffeomorphisms of the covariance group. This can be seen by evaluating the variation of the pseudo-metric under $SU(3)_{color}$ [7]; the homogeneous $SU(3)$ variation δ_ϵ , i.e. $\delta_\epsilon B_\mu^a = \partial_\mu \epsilon^a + B_\mu^b (\lambda_b)_c^a \epsilon^c$, λ_b being the $SU(3)_{color}$ group generators.

For the di-gluon we thus have

$$\delta_\epsilon G_{\mu\nu} = g_{ab}(\partial_\mu \epsilon^a) B_\nu^b + g_{ab} B_\mu^a (\partial_\nu \epsilon^b) + i g_{ab} f_{cd}^a B^c \epsilon^d B_\nu^b + i g_{ab} f_{cd}^b B_\mu^a B_\nu^c \epsilon^d.$$

The homogeneous $SU(3)_{color}$ variation $i g_{ab} f_{cd}^a B^c \epsilon^d B_\nu^b + i g_{ab} f_{cd}^b B_\mu^a B_\nu^c \epsilon^d = i f_{bcd} (B_\mu^b B_\nu^c + B_\mu^c B_\nu^b) \epsilon^d$ vanishes due to total antisymmetry of the $SU(3)_{color}$ (compact group) structure constants. We write now $B_\mu^a = N_\mu^a + A_\mu^a$, and thus

$$\delta_\epsilon G_{\mu\nu} = g_{ab}(\partial_\mu \epsilon^a N_\nu^b + N_\mu^a \partial_\nu \epsilon^b) + g_{ab}(\partial_\mu \epsilon^a A_\nu^b + A_\mu^a \partial_\nu \epsilon^b).$$

As to the terms in A_μ^a, A_ν^b , integration by parts yields $g_{ab}(\epsilon^a \partial_\mu A_\nu^b + \partial_\nu A_\mu^a \epsilon^b)$. But taking Fourier transforms - i.e. the matrix elements for these gluon fluctuations - we find that *these terms are precisely those that vanish in the IR region*. The terms involving the constant N_μ^a, N_ν^b can be rewritten in terms of effective pseudo-diffeomorphisms, defined by

$$\delta_\epsilon G_{\mu\nu} = \partial_\mu \xi_\nu + \partial_\nu \xi_\mu, \quad \xi_\mu \equiv g_{ab} \epsilon^a N_\mu^b. \quad (9)$$

As a result, $G_{\mu\nu}(x)$ transforms as a world tensor, for "artificial" local pseudo-diffeomorphisms ξ_μ . We note that $G_{\mu\nu} = g_{ab} N_\mu^a N_\nu^b + \dots$ is a non-singular (invertible) tensor, thus acting as a metric. It was shown long ago [8] that any spin 2 field will behave and couple like the graviton; it will stay massless because of Lorentz invariance, conservation of the energy-momentum tensor and Einstein Covariance relating to these pseudo-diffeomorphisms. Thus, the local $SU(3)_{color}$ gauge variations contain a subsystem ensuring that the $G_{\mu\nu}$ di-gluon indeed act as a "pseudo-metric" field, precisely emulating gravity. This provides proof and precise limitations for our original conjecture [1,9]. $G_{\mu\nu}(x)$ is a Riemannian pseudo-metric, because it preserves the Lorentz group, so that

$$D_\sigma G_{\mu\nu} = 0, \quad (10)$$

where D_σ is the covariant derivative of the effective pseudo-gravity (the connection will be given by a Christoffel symbol constructed with the metric (3)). The effective action for the IR (zero-color) hadron sector of QCD, written as a pseudo-gravitational theory, with matter in $\overline{SL}(4, R)$ manifolds was derived recently [7] by requiring Riemannian structure defined by the pseudo-metric $G_{\mu\nu}$ even in the presence of the $\overline{SL}(4, R)$ hadronic matter currents. The action reads

$$I = \int d^4x \sqrt{-G} \{ -a R_{\mu\nu} R^{\mu\nu} + b R^2 - c l_G^{-2} R + l_S^{-2} \Sigma_{\alpha\beta}^\gamma \Sigma^{\alpha\beta}_\gamma + l_Q^{-2} \Delta_{\alpha\beta}^\gamma \Delta^{\alpha\beta}_\gamma + L_M \}. \quad (11)$$

The first three terms (R is the scalar curvature and $R_{\mu\nu}$ is the Ricci tensor) constitute the renormalizable quantum GR lagrangian [10]. The fourth and fifth terms in are spin-spin and shear-shear contact interaction terms. The a, b, c are dimensionless constants; l_G, l_S and l_Q , have the dimensions of lengths; from our knowledge of the hadrons we estimate them to be of hadron size $\sim 1 \text{ GeV}$. The pseudo-metric field equations are given by the following expression (; denotes covariant derivative)

$$-a R^{\mu\nu}{}_{;\sigma}{}^{;\sigma} + (a - 2b) R^{;\mu;\nu} - (\frac{1}{2}a - 2b) G^{\mu\nu} R_{;\sigma}{}^{;\sigma} + 2a R^{\mu\sigma\nu\tau} R_{\sigma\tau}$$

$$\begin{aligned}
& -\frac{1}{2}G^{\mu\nu}(aR_{\sigma\tau}R^{\sigma\tau} - bR^2) - 2bRR^{\mu\nu} + cl_G^{-2}(R^{\mu\nu} - \frac{1}{2}G^{\mu\nu}R) \\
& -\frac{1}{2}l_S^{-2}\Sigma_{\alpha\beta}^{\gamma}\Sigma^{\alpha\beta}_{\gamma}G^{\mu\nu} - \frac{1}{2}l_Q^{-2}\Delta_{\alpha\beta}^{\gamma}\Delta^{\alpha\beta}_{\gamma}G^{\mu\nu} = \frac{1}{2}\Theta^{\mu\nu}.
\end{aligned} \tag{12}$$

It is obvious from this expression that $R_{\mu\nu}R^{\mu\nu}$ and R^2 terms in the action yield the fourth-order derivative equation terms, while R yields the usual second-order derivative terms. The latter terms, i.e. r^{-1} -like static solutions, are relevant at "short" distances, while due to a rather soft r -dependence they can be neglected at "large" distances (IR region). Thus, we set $c = 0$ in the following discussion.

LINEAR REGGE TRAJECTORIES

We linearize our theory in terms of $H_{\mu\nu}(x) = G_{\mu\nu}(x) - \eta_{\mu\nu}$, where $\eta_{\mu\nu}$ is the Minkowski metric. The appropriate action, which is bilinear in the $H_{\mu\nu}$ field has the following form

$$\begin{aligned}
I = \int d^4x \{ & \frac{1}{4}aH_{\rho\sigma}\Box^2P_{(2)}^{\rho\sigma\mu\nu}H_{\mu\nu} + (a - 3b)H_{\rho\sigma}\Box^2P_{(0)}^{\rho\sigma\mu\nu}H_{\mu\nu} \\
& - \frac{1}{2}l_S^{-2}H^{\mu\nu}\Sigma_{\alpha\beta}^{\gamma}\Sigma^{\alpha\beta}_{\gamma}H_{\mu\nu} - \frac{1}{2}l_Q^{-2}H^{\mu\nu}\Delta_{\alpha\beta}^{\gamma}\Delta^{\alpha\beta}_{\gamma}H_{\mu\nu} - \frac{1}{2}\Theta^{\mu\nu}H_{\mu\nu} \}.
\end{aligned} \tag{13}$$

The two, respectively $J = 2$ and $J = 0$, completely transverse projectors $P_{(2)}^{\rho\sigma\mu\nu}$ and $P_{(0)}^{\rho\sigma\mu\nu}$ are defined in terms of $\theta^{\mu\nu} = \eta^{\mu\nu} - \partial^\mu\partial^\nu/\Box$,

$$\begin{aligned}
P_{(2)}^{\rho\sigma\mu\nu} &= \frac{1}{2}(\theta^{\rho\mu}\theta^{\sigma\nu} + \theta^{\rho\nu}\theta^{\sigma\mu}) - P_{(0)}^{\rho\sigma\mu\nu} \\
P_{(0)}^{\rho\sigma\mu\nu} &= \frac{1}{3}\theta^{\rho\sigma}\theta^{\mu\nu}
\end{aligned} \tag{14}$$

Raising and lowering of indices is done with the Minkowski metric $\eta_{\mu\nu}$. Note that the Abelian gauge invariance of the linearized action involves the absence of differential operators containing $\partial^\mu\partial^\nu/\Box$. *An important straightforward consequence is the absence of the $J = 1$ part of the $H_{\mu\nu}$ in the action [10].*

We take for simplicity $b = \frac{1}{4}a$ and write, on account of the $J = 1$ mode absence, the bilinear action in the form

$$I = \int d^4x \{ H_{\rho\sigma} [\frac{1}{4}a\Box^2 - \frac{1}{2}(l_S^{-2}\Sigma_{\alpha\beta}^{\gamma}\Sigma^{\alpha\beta}_{\gamma} + l_Q^{-2}\Delta_{\alpha\beta}^{\gamma}\Delta^{\alpha\beta}_{\gamma})] P_{(2+0)}^{\rho\sigma\mu\nu}H_{\mu\nu} - \frac{1}{2}\Theta^{\mu\nu}H_{\mu\nu} \}. \tag{15}$$

The projector $P_{(2+0)}^{\rho\sigma\mu\nu} = P_{(2)}^{\rho\sigma\mu\nu} + P_{(0)}^{\rho\sigma\mu\nu}$ insures that the only physical, propagating modes, are $J^P = 2^+$ and $J^P = 0^+$.

Taking just the homogeneous part, as required for the evaluation of the propagator, we get for the $H_{\mu\nu}$ field the equation of motion

$$[\frac{1}{4}a\Box^2 - \frac{1}{2}l_S^{-2}\Sigma_{\alpha\beta}^{\gamma}\Sigma^{\alpha\beta}_{\gamma} - \frac{1}{2}l_Q^{-2}\Delta_{\alpha\beta}^{\gamma}\Delta^{\alpha\beta}_{\gamma}](P_{(2+0)}H)_{\mu\nu}(x) = 0, \tag{16}$$

which becomes in momentum space

$$[\frac{1}{4}a(p^2)^2 - \frac{1}{2}l_S^{-2}f_S M_\alpha^\beta M_\beta^\alpha - \frac{1}{2}l_Q^{-2}f_Q T_\alpha^\beta T_\beta^\alpha](\tilde{P}_{(2+0)}H)_{\mu\nu}(p) = 0. \tag{17}$$

We have factored out in this expression the $\overline{SL}(4, R)$ group factors (bilinear terms in the algebra's generators). The factors f_S and f_Q represent the residual part of the configuration space integrals, and

$$\tilde{P}_{(2+0)}^{\rho\sigma\mu\nu} = \frac{1}{2}(\tilde{\theta}^{\rho\mu}\tilde{\theta}^{\sigma\nu} + \tilde{\theta}^{\rho\nu}\tilde{\theta}^{\sigma\mu}),$$

where $\tilde{\theta}^{\mu\nu} = \eta^{\mu\nu} - p^\mu p^\nu / p^2$. We write in the following $H_{\mu\nu}$ for $(P_{(2+0)} H_{\mu\nu})$.

For pseudo-gravity, we may regard these equations as *the dynamical quantum equations above the theory's "vacuum"* as represented by hadron matter itself. These equations represent the excitations produced over that ground state by the pseudo-gravity potential, i.e. they are like equations for the $H_{\mu\nu}$ field in an external field of hadronic matter.

We have shown recently [7] that these equations in a rest frame yield that all hadronic states belonging to a single unitary "little-group" irreducible representation lie on a single trajectory in the Chew-Frautschi plane, i.e.

$$(J + \frac{1}{2})^2 = (\alpha' m^2)^2 + \alpha_0^2, \quad (18a)$$

$$(\alpha')^2 = [\frac{2}{a}(l_S^{-2} f_S + l_Q^{-2} f_Q)]^{-1}, \quad (18b)$$

$$\alpha_0^2 = \frac{1}{4} + \frac{l_Q^{-2} f_Q}{l_S^{-2} f_S + l_Q^{-2} f_Q} C_{\overline{SL}(3,R)}^2. \quad (18c)$$

where $C_{\overline{SL}(3,R)}^2$ is the $\overline{SL}(3, R)$ quadratic invariant, and α' is the (asymptotic) trajectory slope. Note that Eq. (18) is meaningful only if $C_{\overline{SL}(3,R)}^2 < 0$. Indeed, this is the case for the (relevant) unitary irreducible $\overline{SL}(3, R)$ representations [11]). Neglecting a slight bending at small m^2 , i.e. the α_0^2 term, we obtain the linear Regge trajectory

$$J = \alpha' m^2 - \frac{1}{2}. \quad (19).$$

Note that for all $\overline{SL}(3, R)$ subrepresentations of a given $\overline{SL}(4, R)$ representation one has the same α' value, i.e.

$$\alpha' = \text{const.} \quad (20)$$

In this manner we have shown that the effective pseudo-gravity theory of the IR QCD regime implies the $\overline{SL}(4, R)$ classification of hadrons as well as, for $\overline{SL}(3, R)$ multiplets with $\Delta J = 2$, the linear Regge trajectory relation which was used in [2,3] as an additional phenomenological input.

It is important to note here that this dynamically derived result could a priori have been in contradiction with some kinematical algebraic constraint relating m^2 to J (such as the notorious Gell-Mann/Dashen "angular condition" upon the representations of the local current algebras). This would result from a study of the Casimir invariants of the $\overline{SA}(4, R)$ group. Should this have been the case, we could have been forced to abandon (19), (20). However, a recent explicit evaluation of the $\overline{SA}(4, R) \supset \overline{SA}(3, R) \supset \overline{SA}(2, R)$ group chain Casimir invariants [12] shows that there are no kinematical m^2 vs. J relations imposed in the specific class of unirreps we use for the particles, i.e. the class in which the "little group" is an artificial $\overline{SA}(3, R)$ whose corresponding unphysical momenta are

put to zero, as is done for the $E(2)$ little group's unphysical momenta for Poincaré group massless particles. The values of the unphysical momenta in $\overline{SA}(3, R)$ enter the $\overline{SA}(4, R)$ Casimir and make it vanish altogether. Thus, our dynamical result Eq. (19) is a bonafide result as well as a prediction of our QCD based theory for the spectrum of hadronic states (organized according to an irreducible $\overline{SL}(3, R)$ subrepresentation of the hadronic matter manifold $\overline{SL}(4, R)$ representation).

Let us consider the static solution of the pseudo-gravity equation in the IR region inside hadronic matter, i.e. for $r \leq 0.8$ fm. This region is dominated by the pure pseudo-gravity terms. The corresponding momentum space Green function is given by

$$G^{\rho\sigma\mu\nu}(p) = \frac{a}{4} \frac{\tilde{P}^{\rho\sigma\mu\nu}_{(2+0)}}{(p^2)^2}, \quad (21)$$

while the pseudo gravity potential $H = H^\mu_\mu$ has in the static case the following form

$$\frac{a}{4}(\nabla^2)^2 H(r) = (\alpha')^{-1} \delta^3(r), \quad (22)$$

for a point source with typical hadronic strength. The solution is a well known linear *confining potential*

$$H(r) = \frac{1}{2\pi a\alpha'} r. \quad (23)$$

HADRON SPECTROSCOPY

When dealing with the hadron Hilbert space states, momenta come in and translations thus have to be adjoined to the algebra. Here we get $sa(4, R) = t_4 \circledast sl(4, R)$ algebra of the $\overline{SA}(4, R) = T_4 \circledast \overline{SL}(4, R)$ group. The massive states of the hadron spectrum are then classified according to the stability subalgebra, here $sa'(3, R) = t'_3 \circledast sl(3, R)$. The t'_3 quantum numbers are trivialized, as is done with the formal translations t'_2 of the Euclidean two-dimensional stability subalgebra, for the massless states in the Poincaré group representations. Hadron states are then characterized by the $sl(3, R)$ subalgebra, whose infinite representations correspond to Regge trajectories. A Regge trajectory described by such a representation corresponds to a given "bag". Spinors are taken care of by the infinite representations of the double covering groups $\overline{SA}(4, R) = T_4 \circledast \overline{SL}(4, R)$ [5]. We collect all hadronic field configurations obtained by successive application of the quantum di-gluon field $sl(4, R)$ dynamical algebra, into an infinite-component field (manifold). These manifolds are subjects to the following constraints: (i) All $SL(4, R)$ representation labels are solely determined by its (enveloping) algebra/subalgebra labels, (ii) owing to $D_\sigma G_{\mu\nu}(x) = 0$, the wave-equation have to be Lorentz covariant, (iii) The lowest manifold components (lowest $\overline{SO}(4)$ subrepresentations) should fit the basic quark system field configuration, i.e. they are fixed by either $\bar{q}q$ or $\bar{q}qA$ color-neutral configurations (not connected by the $SL(4, R)$ shear operators), and (iv) They should transform according to non-unitary representations of the $\overline{SO}(1, 3) \subset \overline{SL}(4, R)$ in order to meet the

experimental fact that a boosted particle keeps its spin quantum number. These natural requirements determine uniquely the selection of manifolds and their equations [2,3,13], while the "non-unitarity" condition on $\overline{SO}(1,3)$ representations is achieved by making use of "A-deunitarized" $\overline{SL}(4,R)$ unitary irreducible representations [5].

If one assumes the baryon wave equation to be of a Dirac-type form

$$(iX^\mu \partial_\mu - M)\Psi(x) = 0, \quad (24)$$

where the infinite matrices X^μ transform as a $(\frac{1}{2}, \frac{1}{2})$ $SL(2, C)$ four-vector representation, one finds [14] a *unique* nontrivial equation of the form (24)

$$\Psi(x) \sim D^{disc}(\frac{1}{2}, 0) \oplus D^{disc}(0, \frac{1}{2}). \quad (25)$$

The wave equation (24) yields a large reduction in the number of states. Thus, in the case of baryons we have a unique choice of the system based on the $[D_{SL(4,R)}^{disc}(\frac{1}{2}, 0) \oplus D_{SL(4,R)}^{disc}(0, \frac{1}{2})]^A$ system. For the decuplet states we have to make use of the symmetrized product of this reducible representation and the finite-dimensional $\overline{SL}(4,R)$ representation $(\frac{1}{2}, \frac{1}{2})$ (generalizing the Rarita - Schwinger approach). The wave equation for the infinite-component Rarita-Schwinger field reads [3]

$$(iX^\mu \partial_\mu - M)\Psi_\rho(x) = 0, \quad (26)$$

while the corresponding manifold transforms as follows

$$\Psi_\rho(x) \sim \{[D^{disc}(\frac{1}{2}, 0)^A \oplus D^{disc}(0, \frac{1}{2})^A] \otimes D^{(1/2, 1/2)}\}_{sym}. \quad (27)$$

The $\overline{SL}(4,R)$ generators have definite space-time properties, and in particular a constrained behavior under the parity operation: The J_i , T_{ij} , and T_{00} operators are parity even, while the K_i and N_i are parity odd. All states of the same $\overline{SL}(3,R)$ subgroup unirrep (Regge trajectory) thus have the same parity; the states of an $SL(2,C)$ or an $\overline{SO}(4)$ subgroup representation have alternating parities. For a given $SL(2,C) = \overline{SO}(4)^A$ representation (j_1, j_2) , the total (spin) angular momentum is $J = J^{(1)} + J^{(2)}$, while the boost operator is given by $K = J^{(1)} - J^{(2)}$. We find the following J^P content of a (j_1, j_2) $\overline{SO}(4)^A$ representation:

$$J^P = (j_1 + j_2)^P, (j_1 + j_2 - 1)^{-P}, (j_1 + j_2 - 2)^P, \dots, (|j_1 - j_2|)^{\pm P}. \quad (28)$$

Thus, by assigning a given parity to any state of an $\overline{SL}(4,R)$ representation, say the lowest state, the parities of all other states are determined.

In the case of mesons, the lowest $SO(4)$ subrepresentations are fixed by either $\overline{q}q$ or $\overline{q}qA$ color-neutral configurations (not connected by the $\overline{SL}(4,R)$ shear operators). The first requirement is met by selecting the $\overline{SL}(4,R)$ multiplicity-free representations, while the third one allows three $SO(4)$ representations: $(0,0)$, $(\frac{1}{2}, \frac{1}{2})$ and $(1,1)$. The $(0,0)$ and $(1,1)$ representations are mutually connected by the shear operators, and thus there are finally two $\overline{SL}(4,R)$ representation candidates for meson manifolds: $D^{ladd}(\frac{1}{2}, \frac{1}{2})^A$

and $D^{ladd}(0,0)^{\mathcal{A}}$. In other words a quark-antiquark colorless manifold transforms w.r.t. the $D^{ladd}(\frac{1}{2}, \frac{1}{2})^{\mathcal{A}} \oplus D^{ladd}(0,0)^{\mathcal{A}}$ $\overline{SL}(4, R)$ representation and couples universally to the pseudo-metric field. The corresponding manifolds $\Phi^{(\frac{1}{2}, \frac{1}{2})}$ and $\Phi^{(0,0)}$ satisfy the following wave equations [3] respectively

$$(\square + M^2)\Phi^{(\frac{1}{2}, \frac{1}{2})} = 0, \quad (\square + M^2)\Phi^{(0,0)} = 0, \quad (29)$$

We have shown recently [13] that adding the P and C parities results in the $SL(4, R) \oplus [Z_2(P) \otimes Z_2(C)]$ classification scheme. Thus, *we assign all hadron states of a given flavor to the wave-equation-projected states corresponding to parity-doubled $\overline{SL}(4, R)^{\mathcal{A}}$ irreducible representations* (their lowest - J states have opposite parities).

$$\text{Baryons } \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array} : [D_{SL(4,R)}^{disc}(\frac{1}{2}, 0) \oplus D_{SL(4,R)}^{disc}(0, \frac{1}{2})]^{\mathcal{A}}, \Psi,$$

$$\{(j_1, j_2)\} = \{(\frac{1}{2}, 0), (\frac{3}{2}, 1), (\frac{5}{2}, 2), \dots\} \oplus \{(0, \frac{1}{2}), (1, \frac{3}{2}), (2, \frac{5}{2}), \dots\}. \quad (30)$$

$$\text{Baryons } \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array} : \{[D_{SL(4,R)}^{disc}(\frac{1}{2}, 0) \oplus D_{SL(4,R)}^{disc}(0, \frac{1}{2})]^{\mathcal{A}} \otimes D^{(\frac{1}{2}, \frac{1}{2})}\}_{sym}, \Psi_{\rho},$$

$$\{(j_1, j_2)\} = \{(1, \frac{1}{2}), (2, \frac{3}{2}), (3, \frac{5}{2}), \dots\} \oplus \{(\frac{1}{2}, 1), (\frac{3}{2}, 2), (\frac{5}{2}, 3), \dots\}. \quad (31)$$

$$\text{Mesons: } [D_{SL(4,R)}^{ladd}(\frac{1}{2}; e_2) \oplus D_{SL(4,R)}^{ladd}(0; e_2)]^{\mathcal{A}}, \Phi,$$

$$\{(j_1, j_2)\} = \{(\frac{1}{2}, \frac{1}{2}), (\frac{3}{2}, \frac{3}{2}), (\frac{5}{2}, \frac{5}{2}), \dots\} \oplus \{(0, 0), (1, 1), (2, 2)\}. \quad (32)$$

The Young tableaux in (30) and (31), correspond to the flavor - $SU(6)$ assignments. The $\overline{SO}(4)^{\mathcal{A}}$ states of (30) and (31), when reorganized with respect to the $\overline{SL}(3, R)$ subgroup, form an infinite sum of Regge-type $\Delta J = 2$ recurrences with the J content $\{J\} = \{\frac{1}{2}, \frac{5}{2}, \frac{9}{2}, \dots\}, \{J\} = \{\frac{3}{2}, \frac{7}{2}, \frac{11}{2}, \dots\}$.

Owing to a presence of non trivial X matrix in the baryon manifold wave equation, we take for baryons the following mass formula

$$m^2 = m_0^2 + (\alpha')^{-1}(j_1 + j_2 - \frac{1}{2} - \frac{1}{2}n), \quad (33)$$

where m_0 is the mass of the lowest-lying state and $n = j_1 + j_2 - J$. We predict from the linear Regge formula, for the $SO(4)$ -degenerate meson recurrences, the mass spectrum given by

$$m_{(\frac{1}{2}, \frac{1}{2})}^2 : m_{(\frac{3}{2}, \frac{3}{2})}^2 : m_{(\frac{5}{2}, \frac{5}{2})}^2 : \dots = 1 \cdot (\alpha')^{-1} : 2 \cdot (\alpha')^{-1} : 3 \cdot (\alpha')^{-1} : \dots. \quad (34)$$

i.e. for $(\alpha')^{-1} \approx 1 \text{ GeV}^2$ we have $m_{(\frac{1}{2}, \frac{1}{2})} : m_{(\frac{3}{2}, \frac{3}{2})} : m_{(\frac{5}{2}, \frac{5}{2})} : \dots = 1 : 1.41 : 1.73 : \dots$. One finds that $m_{(\frac{1}{2}, \frac{1}{2})}^2 \leq 0.9 \text{ GeV}^2$, and thus $m_{(0,0)}^2 < 0$, i.e. that the $(0,0)$ $\overline{SO}(4)$ multiplet is not realized. Moreover, we find that pairs of $SL(3, R)$ Regge trajectories with opposite PC -parities are degenerate in the Chew - Frautschi plot, i.e. we predict the "exchange degenerate" Regge trajectories.

We organize our predicted $\overline{SL}(4, R)^{\mathcal{A}}$ states (30), (31) and (32) according to the Regge mass formula in Tables I - IV, and confront them with the known $SU(3)$ hadron resonances [15]. We find a good fit with the experimental data.

REFERENCES

1. Dj. Šijački and Y. Ne'eman, *Phys. Lett. B* 247 (1990) 571.
2. Y. Ne'eman and Dj. Šijački, *Phys. Lett. B* 157 (1985) 267.
3. Y. Ne'eman and Dj. Šijački, *Phys. Rev. D* 37 (1988) 3267.
4. Y. Ne'eman and Dj. Šijački, *Phys. Lett. B* 157 (1985) 275.
5. Y. Ne'eman and Dj. Šijački, *Int. J. Mod. Phys. A* 2 (1987) 1655.
6. A. Arima and F. Iachello, *Phys. Rev. Lett.* 35 (1975) 1069.
7. Y. Ne'eman and Dj. Šijački, *Phys. Lett. B* 276 (1992) 173.
8. S. Weinberg, *Phys. Rev. B* 138 (1965) 988.
9. Dj. Šijački and Y. Ne'eman, *Phys. Lett. B* 250 (1990) 1.
10. K. S. Stelle, *Phys. Rev. D* 16 (1977) 953; *Gen. Relat. Grav.* 9 (1978) 353.
11. Dj. Šijački, *J. Math. Phys.* 16 (1975) 298.
12. J. Lemke, Y. Ne'eman and J. Pecina-Cruz, *J. Math. Phys.* 33 (1992) 2656.
13. Dj. Šijački and Y. Ne'eman, Hadrons in an $\overline{SL}(4, R)$ Classification II. Mesons and C, P Assignments, (to be published).
14. A. Cant and Y. Ne'eman, *J. Math. Phys.* 26 (1985) 3180.
15. Particle Data Group, *Phys. Rev. D* 45 S1 (1992).

TABLE I Assignment of N , Λ and Σ $SU(3)$ - octet states.*

$[D(\frac{1}{2}, 0) \oplus D(0, \frac{1}{2})]^{(+)}$					$[D(\frac{1}{2}, 0) \oplus D(0, \frac{1}{2})]^{(-)}$				
(j_1, j_2)	J^P	$\{N\}$	$\{\Lambda\}$	$\{\Sigma\}$	(j_1, j_2)	J^P	$\{N\}$	$\{\Lambda\}$	$\{\Sigma\}$
$(\frac{1}{2}, 0)$	$\frac{1}{2}^+$	$N(940)$	$\Lambda(1116)$	$\Sigma(1193)$	$(0, \frac{1}{2})$	$\frac{1}{2}^-$	$N(1535)$	$\Lambda(1670)$	$\underline{\Sigma}(\sim 1500)$
$(\frac{3}{2}, 1)$	$\frac{1}{2}^+$	$N(1440)$	$\Lambda(1600)$	$\Sigma(1660)$	$(1, \frac{3}{2})$	$\frac{1}{2}^-$	$N(1650)$	$\Lambda(1800)$	$\Sigma(1620)$
	$\frac{3}{2}^-$	$N(1520)$	$\Lambda(1690)$	$\Sigma(1670)$		$\frac{3}{2}^+$	$\underline{N}(1540)$	$\Lambda(1890)$	$\Sigma(1670)$
	$\frac{5}{2}^+$	$N(1680)$	$\Lambda(1820)$	$\Sigma(1915)$		$\frac{5}{2}^-$	$N(1675)$	$\Lambda(1830)$	$\Sigma(1775)$
$(\frac{5}{2}, 2)$	$\frac{1}{2}^+$	$N(1710)$	$\Lambda(1800)$	$\Sigma(1880)$	$(2, \frac{5}{2})$	$\frac{1}{2}^-$	$N(2090)$	$\underline{\Lambda}(2020)$	$\Sigma(1750)$
	$\frac{3}{2}^-$	$N(1700)$	$\Lambda(2000)$	$\Sigma(1840)$		$\frac{3}{2}^+$	$N(1720)$		
	$\frac{5}{2}^+$	$N(2000)$	$\Lambda(2110)$			$\frac{5}{2}^-$	$N(2200)$		
	$\frac{7}{2}^-$	$N(2190)$	$\Lambda(2350)$			$\frac{7}{2}^+$	$N(1990)$		
	$\frac{9}{2}^+$	$N(2220)$				$\frac{9}{2}^-$	$N(2250)$		
$(\frac{7}{2}, 3)$	$\frac{1}{2}^+$	$\underline{N}(2100)$	$\Lambda(2325)$	$\Sigma(2250)$	$(3, \frac{7}{2})$	$\frac{1}{2}^-$			
	$\frac{3}{2}^-$	$N(2080)$				$\frac{3}{2}^+$			
	$\frac{5}{2}^+$					$\frac{5}{2}^-$			
	$\frac{7}{2}^-$					$\frac{7}{2}^+$			
	$\frac{9}{2}^+$					$\frac{9}{2}^-$			
	$\frac{11}{2}^-$		$N(2600)$	$\frac{11}{2}^+$					
	$\frac{13}{2}^+$	$N(2700)$	$\frac{13}{2}^-$						

TABLE II Assignment of Δ and Σ $SU(3)$ - decuplet states.*

$[D(\frac{1}{2}, 0)_\mu \oplus D(0, \frac{1}{2})_\mu]^{(-)}$				$[D(\frac{1}{2}, 0)_\mu \oplus D(0, \frac{1}{2})_\mu]^{(+)}$			
(j_1, j_2)	J^P	$\{\Delta\}$	$\{\Sigma\}$	(j_1, j_2)	J^P	$\{\Delta\}$	$\{\Sigma\}$
$(1, \frac{1}{2})$	$\frac{1}{2}^-$	$\Delta(1620)$		$(\frac{1}{2}, 1)$	$\frac{1}{2}^+$	$\underline{\Delta}(1550)$	$\Sigma(1770)$
	$\frac{3}{2}^+$	$\Delta(1232)$	$\Sigma(1385)$		$\frac{3}{2}^-$	$\Delta(1700)$	$\Sigma(1580)$
$(2, \frac{3}{2})$	$\frac{1}{2}^-$	$\Delta(1900)$	$\underline{\Sigma}(2000)$	$(\frac{3}{2}, 2)$	$\frac{1}{2}^+$	$\Delta(1910)$	$\Sigma(1940)$
	$\frac{3}{2}^+$	$\Delta(1600)$	$\Sigma(1690)$		$\frac{3}{2}^-$	$\underline{\Delta}(1940)$	
	$\frac{5}{2}^-$				$\frac{5}{2}^+$	$\Delta(1905)$	
	$\frac{7}{2}^+$	$\Delta(1950)$	$\Sigma(2030)$		$\frac{7}{2}^-$		
$(3, \frac{5}{2})$	$\frac{1}{2}^-$	$\underline{\Delta}(2150)$	$\Sigma(2080)$	$(\frac{5}{2}, 3)$	$\frac{1}{2}^+$		$\underline{\Sigma}(2070)$ $\underline{\Sigma}(2150)$
	$\frac{3}{2}^+$	$\Delta(1920)$			$\frac{3}{2}^-$		
	$\frac{5}{2}^-$	$\Delta(1930)$			$\frac{5}{2}^+$	$\Delta(2000)$	
	$\frac{7}{2}^+$		$\frac{7}{2}^-$		$\underline{\Delta}(2200)$		
	$\frac{9}{2}^-$	$\Delta(2400)$	$\frac{9}{2}^+$		$\Delta(2300)$		
	$\frac{11}{2}^+$	$\Delta(2420)$	$\Sigma(2620)$		$\frac{11}{2}^-$		
$(4, \frac{7}{2})$	$\frac{1}{2}^-$			$(\frac{7}{2}, 4)$	$\frac{1}{2}^+$		
	$\frac{3}{2}^+$				$\frac{3}{2}^-$		
	$\frac{5}{2}^-$	$\underline{\Delta}(2350)$			$\frac{5}{2}^-$		
	$\frac{7}{2}^+$	$\Delta(2390)$			$\frac{7}{2}^-$		
	$\frac{9}{2}^-$				$\frac{9}{2}^+$		
	$\frac{11}{2}^+$				$\frac{11}{2}^-$		
	$\frac{13}{2}^-$	$\Delta(2750)$			$\frac{13}{2}^+$		
	$\frac{15}{2}^+$	$\Delta(2950)$			$\frac{15}{2}^-$		

* Underlined entries are uncertain.

TABLE III $D^{ladd}(0,0) \oplus D^{ladd}(\frac{1}{2}, \frac{1}{2})$, $J_{min}^P = 0^+$. Assignment of $SU(3)$ Meson States.*

(j_1, j_2)	J^{PC}	$I = 1$	$I = \frac{1}{2}$	$I = 0$	(j_1, j_2)	J^{PC}	$I = 1$	$I = \frac{1}{2}$	$I = 0$
$(\frac{1}{2}, \frac{1}{2})$	0^{++} 1^{--}	$a_0(980)$ $\rho(770)$	$K^*(892)$	$f_0(975)$ $f_0(1400)$ $\omega(783)$ $\phi(1020)$	$(\frac{1}{2}, \frac{1}{2})$	0^{+-} 1^{-+}			
$(1, 1)$	0^{++} 1^{--} 2^{++}	$\underline{a}_0(1320)$ $\rho(1450)$ $a_2(1320)$	$K_0^*(1430)$ $K^*(1370)$ $K_2^*(1430)$	$\underline{f}_0(1240)$ $f_0(1590)$ $\omega(1390)$ $f_2'(1525)$ $f_2(1270)$	$(1, 1)$	0^{+-} 1^{-+} 2^{+-}	$\underline{\hat{\rho}}(1405)$		
$(\frac{3}{2}, \frac{3}{2})$	0^{++} 1^{--} 2^{++} 3^{--}	$\rho(1700)$ $\rho_3(1690)$	$K^*(1680)$ $K_3^*(1780)$	$f_0(1525)$ $f_0(1750)$ $\omega(1600)$ $\phi(1680)$ $\underline{f}_2(1810)$ $\omega_3(1670)$ $\phi_3(1850)$	$(\frac{3}{2}, \frac{3}{2})$	0^{+-} 1^{-+} 2^{+-} 3^{-+}			
$(2, 2)$	0^{++} 1^{--} 2^{++} 3^{--} 4^{++}	$\underline{e}^+ \underline{e}^-(1830)$ $\underline{a}_4(2040)$	$\underline{K}_0^*(1950)$ $\underline{K}_2^*(1980)$ $K_4^*(2045)$	$\underline{f}_0(1750)$ $f_4(2050)$ $\underline{f}_4(2020)$	$(2, 2)$	0^{+-} 1^{-+} 2^{+-} 3^{-+} 4^{+-}			$\underline{X}(1910)$
$(\frac{5}{2}, \frac{5}{2})$	0^{++} 1^{--} 2^{++} 3^{--} 4^{++} 5^{--}	$\underline{\rho}(2150)$ $\underline{\rho}_3(2250)$ $\underline{\rho}_5(2350)$	 $\underline{K}_5^*(2380)$	 $\underline{f}_4(2300)$	$(\frac{9}{2}, \frac{9}{2})$	0^{+-} 1^{-+} 2^{+-} 3^{-+} 4^{+-} 5^{-+}			

TABLE IV $D^{ladd}(0,0) \oplus D^{ladd}(\frac{1}{2}, \frac{1}{2})$, $J_{min}^P = 0^-$. Assignment of $SU(3)$ Meson States.*

(j_1, j_2)	J^{PC}	$I = 1$	$I = \frac{1}{2}$	$I = 0$	(j_1, j_2)	J^{PC}	$I = 1$	$I = \frac{1}{2}$	$I = 0$
$(\frac{1}{2}, \frac{1}{2})$	0^{-+} 1^{+-}	$\pi(140)$ $b_1(1235)$	$K(494)$ $K_1(1270)$	$\eta(549)\eta'(958)$ $h_1(1170)$	$(\frac{1}{2}, \frac{1}{2})$	0^{--} 1^{++}	$a_1(1260)$	$K_1(1400)$	$f_1(1285)f_1(1420)$
$(1, 1)$	0^{-+} 1^{+-} 2^{-+}	$\pi(1300)$	$\underline{K}(1460)$ $\underline{K}_1(1650)$ $\underline{K}_2(1580)$	$\eta(1295)\eta(1440)$ $\underline{h}_1(1380)$	$(1, 1)$	0^{--} 1^{++} 2^{--}			$f_1(1510)$
$(\frac{3}{2}, \frac{3}{2})$	0^{-+} 1^{+-} 2^{-+} 3^{+-}	$\underline{\pi}(1770)$ $\pi_2(1670)$	$\underline{K}(1830)$ $K_2(1770)$		$(\frac{3}{2}, \frac{3}{2})$	0^{--} 1^{++} 2^{--} 3^{++}			
$(2, 2)$	0^{-+} 1^{+-} 2^{-+} 3^{+-} 4^{-+}				$(2, 2)$	0^{--} 1^{++} 2^{--} 3^{++} 4^{--}	$\underline{a}_3(2050)$		
$(\frac{5}{2}, \frac{5}{2})$	0^{-+} 1^{+-} 2^{-+} 3^{+-} 4^{-+} 5^{+-}	$\pi_2(2100)$	$\underline{K}_2(2250)$ $\underline{K}_3(2320)$ $\underline{K}_4(2500)$	$\underline{\eta}(2100)$	$(\frac{5}{2}, \frac{5}{2})$	0^{--} 1^{++} 2^{--} 3^{++} 4^{--} 5^{++}			

* Underlined entries are uncertain.